

The flight of the bumblebee: solutions from a vector-induced spontaneous Lorentz symmetry breaking model

Orfeu Bertolami* and Jorge Páramos*

**Instituto Superior Técnico, Departamento de Física,
Av. Rovisco Pais 1, 1049-001 Lisboa, Portugal*

Abstract. The vacuum solutions arising from a spontaneous breaking of Lorentz symmetry due to the acquisition of a vacuum expectation value by a vector field are derived. These include the purely radial Lorentz symmetry breaking (LSB), radial/temporal LSB and axial/temporal LSB scenarios. It is found that the purely radial LSB case gives rise to new black hole solutions. Whenever possible, Parametrized Post-Newtonian (PPN) parameters are computed and compared to observational bounds, in order to constrain the Lorentz symmetry breaking scale.

1. INTRODUCTION

Lorentz invariance is clearly one of the most fundamental symmetries of Nature. It is both theoretically sound and experimentally well tested [1, 2], thus playing a leading role in most theories of gravity. Therefore, it is only natural that little attention has been paid to the consequences of explicitly breaking this symmetry.

A more flexible approach to this question admits a spontaneous breaking of this symmetry, instead of an explicit one, analogously to the Higgs mechanism in the Standard Model of particle physics [3, 4, 5, 6]. This can arise if a vector field ruled by a potential exhibiting a minimum rolls to its vacuum expectation value (vev) – this vector field, usually referred to as “bumblebee” vector, thus acquires a specific four-dimensional orientation.

From a theoretical standpoint, a spontaneous Lorentz symmetry breaking (LSB) is possible, for instance, in string/M-theory, arising from non-trivial solutions in string field theory [3, 4] and in noncommutative field theories [7, 8]. A spacetime variation of fundamental coupling constants could also lead to a spontaneous LSB [9]. Experimentally, the violation of Lorentz invariance could be tested in ultra-high energy cosmic rays [10].

The consequences of the “bumblebee” vector scenario were studied in Ref. [11]; in there, three relevant cases were taken into account: the bumblebee field acquiring a purely radial vev, a mixed radial and temporal vev and a mixed axial and temporal vev. The results were analyzed in terms of the PPN parameters, when possible, prompting for comparison with current and future experimental bounds and effects, for instance, from string theory in a low-energy scenario. These bounds may arise from the observations of the Bepi-Colombo [12] and LATOR [13] missions (see also Ref. [14]) for a discussion on future gravitational experiments).

The action of the bumblebee model is written as

$$S = \int d^4x \sqrt{-g} \left[\frac{1}{2\kappa} (R + \xi B^\mu B^\nu R_{\mu\nu}) - \frac{1}{4} B^{\mu\nu} B_{\mu\nu} - V(B^\mu B_\mu \pm b^2) \right], \quad (1)$$

where $\kappa = 8\pi G$, $B_{\mu\nu} = \partial_\mu B_\nu - \partial_\nu B_\mu$, ξ is a coupling constant and b^2 sets the bumblebee’s vev, since the potential V driving Lorentz and/or CPT violation is supposed to have a minimum at $B^\mu B_\mu \pm b^2 = 0$, i.e. $V'(b_\mu b^\mu) = 0$. The particular form of this potential is irrelevant, since one assumes that the bumblebee field is frozen at its vev. The scale of b_μ should be obtained from string theory or from a low-energy extension to the Standard Model. Hence, one expects b_μ to be of order of M_{Pl} , the Planck mass, or M_{EW} , the electroweak breaking scale.

2. PURELY RADIAL LSB

In this section, a method to obtain the exact solution for the purely radial LSB is developed. A static, spherically symmetric spacetime, with a Birkhoff metric $g_{\mu\nu} = \text{diag}(-e^{2\phi}, e^{2\rho}, r^2, r^2 \sin^2 \theta)$ is considered. It can be easily seen that the Killing vectors of the metric are conserved, showing that radial symmetry is still valid; this enables the construction of a covariantly conserved current associated with the vector bumblebee field [11].

The affine connection derived from the metric $g_{\mu\nu}$ allows for the computation of b_μ , given the non-trivial covariant derivative with respect to the radial coordinate, and taking $b_\mu = (0, b(r), 0, 0)$. Hence, from $D_\mu b_\nu = \partial_\mu b_\nu - \Gamma_{\nu\mu}^\alpha b_\alpha = 0$, it follows that $b(r) = \xi^{-1/2} b_0 e^\rho$, where the factor $\sqrt{\xi}$ is introduced for later convenience. As expected, $b^2 = b^\mu b_\mu = b_0^2 \xi^{-1}$ is constant.

The (spatial) action can be thus written as

$$S_s = \int d^4x \sqrt{-g} \left[\frac{R}{2\kappa} + (g^{rr})^2 b^2(r) R_{rr} \right]_s. \quad (2)$$

The determinant of the metric is given by $\sqrt{-g} = r^2 e^{\rho+\phi}$; the scalar curvature and the relevant non-vanishing Ricci tensor component are given by $R = 2r^{-2} [1 + (2r\rho' - 1)e^{-2\rho}]$ and $R_{rr} = 2r^{-1}\rho'$, where the prime stands for derivative with respect to r and we have integrated over the angular dependence. Also, $\xi (b^r)^2 R_{rr} = b_0^2 2r^{-1} \rho' e^{-2\rho}$, where b^r is the contravariant radial component of b_μ . By introducing the field redefinition $\Psi = (1 - e^{-2\rho}) r^{-2}$, the action may be rewritten as [15]

$$S_s = \frac{2}{\kappa} \int dr e^{\rho+\phi} r^2 \left[(3 + b_0^2) \Psi + (1 + \frac{b_0^2}{2}) r \Psi' \right]. \quad (3)$$

Variation with respect to ϕ produces the equation of motion

$$(3 + b_0^2) \Psi + (1 + \frac{b_0^2}{2}) r \Psi' = 0, \quad (4)$$

which admits the solution $\Psi(r) = \Psi_0 r^{-3+L}$, with $L \equiv 3 - (3 + b_0^2)/(1 + b_0^2/2) \simeq b_0^2/2$, and hence $L \simeq b_0^2/2$. We thus obtain $g_{rr} = e^{2\rho} = (1 - \Psi_0 r^{-1+L})^{-1}$. Comparison with the usual Schwarzschild metric yields

$$g_{rr} = \left(1 - \frac{2G_L m}{r} r^L \right)^{-1}, \quad g_{00} = -1 + \frac{2G_L m}{r} r^L, \quad (5)$$

where G_L has dimensions $[G_L] = L^{2-L}$ (in natural units, where $c = \hbar = 1$); one can define $G_L = G r_0^{-L}$, where r_0 is an arbitrary distance. The $L \rightarrow 0$ limit yields $G_L \rightarrow G$ and the usual geometrical mass, $M \equiv Gm$, with dimensions of length. From now on we express all results in terms of M . The event horizon condition is given by

$$g_{00} = -1 + \frac{2M}{r} \frac{r^L}{r_0} = 0, \quad (6)$$

thus $r_s = (2M r_0^{-L})^{1/(1-L)}$. The norm-square of the Riemann tensor, $I = R_{\mu\nu\rho\lambda} R^{\mu\nu\rho\lambda}$, is found to be

$$I = 48 \left[1 - \frac{5}{3}L + \frac{17}{12}L^2 - \frac{1}{2}L^3 + \frac{1}{12}L^4 \right] M^2 \left(\frac{r}{r_0} \right)^{2L} r^{-6} \simeq \left(1 - \frac{5}{3}L \right) \left(\frac{r}{r_0} \right)^{2L} I_0, \quad (7)$$

where $I_0 = 48M^2 r^{-6}$ is the usual scalar invariant in the limit $L \rightarrow 0$. Since $I(r = r_s)$ is finite, the singularity at $r = r_s$ is removable; accordingly, the singularity at $r = 0$ is intrinsic, as the scalar invariant diverges there. One concludes that an axial LSB gravity model admits new black hole solutions with a singularity well protected within a horizon of radius r_s . The associated Hawking temperature is

$$T = \frac{\hbar}{k_B} \frac{1}{4\pi r_s} = (2M r_0^{-L})^{-L/(1-L)} T_0 \simeq (2M r_0^{-L})^{-L} T_0, \quad (8)$$

where $T_0 = \hbar/8\pi k_B M$ is the usual Hawking temperature, recovered in the limit $L \rightarrow 0$.

Since the obtained metric cannot be expanded in powers of $U = M/r$, a PPN expansion is not feasible. However, a comparison with results for deviations from Newtonian gravity [16], usually stated in terms of a Yukawa potential of the form

$$V_Y(r) = \frac{G_Y m}{r} \left(1 + \alpha e^{-r/\lambda}\right), \quad (9)$$

yields $G_L r^L = G_Y (1 + \alpha e^{-r/\lambda})$, which, to first order around $r = r_0$, reads

$$G_L r_0^L \left(1 + L \frac{r}{r_0}\right) = G_Y \left(1 + \alpha - \alpha \frac{r}{\lambda}\right), \quad (10)$$

so that one identifies $\lambda = r_0$ and $\alpha = -L$ (with $G_Y(1 - L) = G_L r_0^L = G$). Planetary tests to Kepler's law in Venus indicate that $\lambda = r_0 = 0.723 \text{ AU}$ and $L = |\alpha| \leq 2 \times 10^{-9}$.

3. RADIAL/TEMPORAL LSB

We consider now the mixed radial and temporal Lorentz symmetry breaking. As before, it is assumed that the bumblebee field B_μ has relaxed to its vacuum expectation value. Provided that one takes temporal variations to be of the order of the age of the Universe H_0^{-1} , where H_0 is the Hubble constant, a Birkhoff static, radially symmetric metric $g_{\mu\nu} = \text{diag}(-e^{2\phi}, e^{2\rho}, r^2, r^2 \sin^2(\theta))$ may still be used. The physical gauge choice of a vanishing covariant derivative of the field B_μ yields $b_r(r) = \xi^{-1/2} A_r e^\rho$ and, similarly, $b_0(r) = \xi^{-1/2} A_0 e^\phi$, with A_0 and A_r dimensionless constants. As before, $b^2 = b^\mu b_\mu = (A_r^2 - A_0^2) \xi^{-1}$ is constant.

In the present case, the symmetry $\phi = -\rho$ does not hold, as now both a radial and a temporal component for the vector field v_μ are present; for this reason, the previous spatial action formalism depicted in Eq. (3) cannot be used. Instead, the full Einstein equations must be dealt with,

$$G_{\mu\nu} = \xi \left[\frac{1}{2} (b^\alpha)^2 R_{\alpha\alpha} g_{\mu\nu} - b_\mu b^\nu R_{\mu\nu} - b_\nu b^\mu R_{\nu\mu} \right]. \quad (11)$$

Since the bumblebee field has relaxed to its v_{ev} and therefore both the field strength and the potential term vanish, the additional equation of motion for the vector field is trivial. The metric Ansatz and the expressions for b_μ then yield

$$G_{00} = \frac{1}{2} [3A_0^2 R_{00} - A_r^2 e^{2(\phi-\rho)} R_{rr}] \quad , \quad G_{rr} = \frac{1}{2} [A_0^2 e^{2(\rho-\phi)} R_{00} - 3A_r^2 R_{rr}] \quad , \quad (12)$$

Writing $G_{00} = g_0(r) e^{2(\phi-\rho)}$, $G_{rr} = g_r(r)$, $R_{00} = f_0(r) e^{2(\phi-\rho)}$ and $R_{rr} = f_r(r)$, where

$$\begin{aligned} f_0(r) &\equiv \frac{(2 - r\rho')\phi'}{r} + \phi'^2 + \phi'' \quad , \quad f_r(r) \equiv \frac{(2 + r\phi')\rho'}{r} - \phi'^2 - \phi'' \quad , \\ g_0(r) &\equiv \frac{-1 + e^{2\rho}}{r^2} + \frac{2\rho'}{r} \quad , \quad g_r(r) \equiv \frac{1 - e^{2\rho}}{r^2} + \frac{2\phi'}{r} \quad . \end{aligned}$$

the Einstein equations read

$$g_0(r) = \frac{1}{2} [3A_0^2 f_0(r) - A_r^2 f_r(r)] \quad , \quad g_r(r) = \frac{1}{2} [A_0^2 f_0(r) - 3A_r^2 f_r(r)] \quad . \quad (13)$$

This is the set of coupled second order differential equations which must be solved, with boundary conditions given by $\phi(\infty) = \rho(\infty) = \phi'(\infty) = \rho'(\infty) = 0$.

The spontaneous LSB is clearly exhibited; as can be noticed from $g_0 + g_r = f_0 + f_r$, one has $(1 - 2A_0^2) f_0 = -(1 + 2A_r^2) f_r$; in the unperturbed case $A_0 = A_r = 0$, $f_0 = -f_r$, and the Schwarzschild solution $\phi = -\rho$ is recovered from $g_0 + g_r = 0$. This symmetry does not hold in the perturbed case, which produces $f_0 \approx -(1 + 2A_0^2 + 2A_r^2) f_r$.

An expansion of the metric in terms of $\phi = \phi_0 + \delta\phi$ and $\rho = -\phi_0 - \delta\rho$ allows for the solving of Eqs. (13), where ϕ_0 is given by the usual Schwarzschild metric, $\phi_0 = \frac{1}{2} \ln(1 - 2M/r)$, and $\delta\rho$, $\delta\phi$ are assumed to be small perturbations. After some algebra, the solution is found to be [11]

$$\delta\phi = Kr^{-\alpha} \ , \ \delta\rho \simeq \left[1 + \alpha \frac{(A+B)}{2}\right] Kr^{-\alpha} \ , \quad (14)$$

where $A = A_0^2$, $B = A_r^2$, K is an integration constant and

$$\alpha = \frac{-C_1 - C_2 + \sqrt{(C_1 + C_2)^2 + 4C_1C_3}}{2C_1} > 0 \ , \quad (15)$$

with

$$C_1 = A + 3B + AB + 9B^2 \simeq A + 3B \ , \ C_2 = 2 + B - 3A + 16AB \simeq 2 \ , \ C_3 = 2 + B - 7A \simeq 2 \ . \quad (16)$$

One can linearize the exponent α around $C_1 \ll 1$, yielding $\alpha \simeq (2 - 7A + B)/(2 + A + 3B)$, so that $\alpha \simeq 1$.

After solving the coupled differential Eqs. (13), the non-trivial components of the metric now read

$$\begin{aligned} g_{tt} &= -e^{2(\phi_0 + \delta\phi)} = -e^{2Kr^{-\alpha}} \left(1 - \frac{2M}{r}\right) \ , \\ g_{rr} &= e^{-2(\phi_0 + \delta\rho)} = \frac{e^{-(2 + \alpha(A+B))Kr^{-\alpha}}}{1 - \frac{2M}{r}} \equiv \frac{e^{-2K_r r^{-\alpha}}}{1 - \frac{2M}{r}} \ , \end{aligned} \quad (17)$$

with the definition $K_r \equiv [1 + \alpha(A+B)/2]K \simeq [1 + (A+B)/2]K$. Following the algebra of a Lorentz transformation to a isotropic coordinate system, on which all spatial metric components are equal, and then to a quasi-cartesian referential, the resulting metric is [11]

$$\eta_{tt} = g_{tt} = -1 + 2U - \left(1 - \frac{K + K_r}{M}\right) U^2 \ , \ \eta_{\xi'\xi'} = 1 + \left(1 - \frac{K + 2K_r}{M}\right) U \ , \quad (18)$$

and the PPN parameters may be directly read, yielding $\beta \simeq 1 - (K + K_r)/M$ and $\gamma \simeq 1 - (K + 2K_r)/M$. Inverting this relation gives $K/M \simeq 1 - 2\beta + \gamma$, $K_r/M \simeq \beta - \gamma$. Hence, a temporal/radial LSB manifests itself linearly on the PPN parameters β and γ . A caveat of these results is the clear dependence of the obtained PPN parameters on the free-valued integration constants K and K_r , instead of the physical parameters associated with the breaking of Lorentz invariance. This reflects the linearization procedure followed in order to obtain the radially symmetric Birkhoff metric solution to the Einstein equations.

The bounds derived from the Nordvedt effect, $|\beta - 1| \leq 6 \times 10^{-4}$ [17] and the Cassini-Huygens experiment, $\gamma = 1 + (2.1 \pm 2.3) \times 10^{-5}$ [18], can be used to obtain $|K + K_r| < 0.9 \ m$ and $K + 2K_r = (-3.1 \pm 3.4) \times 10^{-2} \ m$. Since, by definition

$$K_r = \left[1 + \alpha \frac{(A+B)}{2}\right] K \ , \quad (19)$$

with $\alpha \simeq 1$, $A, B \ll 1$, deviations of K_r from K are expected to be small. Thus, considering for instance the constraint $|1 - K_r/K| < 0.1$, one gets $\alpha(A+B) < 0.2$; the limiting case $K \sim K_r$ gives $K \simeq (-1 \pm 1.1) \times 10^{-2}$, indicating a perturbation with a very short range (actually, well inside the Sun, so that one should work with the interior Schwarzschild solution). The range of allowed values for these parameters is depicted in Figures 1 and 2.

In the limit $M \rightarrow 0$, Eqs. (18) yield $\bar{g}_{tt} = -e^{2K/\xi}$ and $\bar{g}_{rr} = -e^{2K_r/\xi}$. An analogy with Rosen's bimetric theory allows for the PPN parameter α_2 to be obtained, by interpreting this change of the metric component as due to a background metric $\eta_{\mu\nu}$ [19]. Notice, however, that the vector field no longer rolls to a radial vev in the absence of a central mass, since this spatial symmetry is inherited from its presence, so this result should be taken with caution; this said, one obtains [20] $\alpha_2 = (c/c_G) - 1 = e^{2(K_r - K)/\xi} - 1 \simeq 2(K_r - K)/\xi = (A+B)\alpha/\xi$, which has a radial dependency. Assuming $\alpha \simeq 1$ and considering the spin precession constraint arising from solar to ecliptic alignment measurements [17], one has $|\alpha_2| = |A+B|/r < 4 \times 10^{-7}$, implying that $|A+B| < 4 \times 10^{-7} r_\odot = 2.78 \times 10^2 \ m$, where $r_\odot = 6.96 \times 10^8 \ m$ is the radius of the Sun.

Further pursuing this analogy, we remark that, since there is no explicit Lorentz breaking, the speed of light remains equal to c . However, the speed of gravitational waves c_G is shifted by an amount




FIGURE 1. Allowed values for K and K_r [11].




FIGURE 2. Detail of Fig. 1, showing only the allowed region [11].

$$\sqrt{\frac{c_G}{c}} = e^{2(K-K_r)/\xi} \simeq \left[1 - \frac{A+B}{\xi} \right], \quad (20)$$

and hence it acquires a radial dependence. As stated before, this result is highly simplistic and should be taken with caution, since it lacks a complete treatment of gravitational radiation induced by LSB, taking into account variations of the bumblebee field B_μ around its vev b_μ .

Finally, notice that, since the radial LSB effects are exact, while the radial/temporal results are not, a direct comparison of these scenarios by taking the $A \rightarrow 0$ limit is not possible.

4. AXIAL/TEMPORAL LSB

The anisotropic LSB case is dealt with in this section. As before, we assume that the bumblebee field is stabilized at its vacuum expectation value, which possesses both a temporal and a spatial component; the latter is taken to lie on the x-axis, that is, $b_\mu = \kappa^{-1}(a, b, 0, 0)$. Since the radial symmetry of the Schwarzschild is clearly broken, one cannot resort to a Birkhoff canonical *Ansatz*. Instead, the perturbations $h_{\mu\nu}$ to the flat Minkowsky metric must be obtained. To first order in $h_{\mu\nu}$, one has

$$\begin{aligned} R_{00} &= -\frac{1}{2}\nabla^2 h_{00} \quad , \quad R_{0i} = -\frac{1}{2}(\nabla^2 h_{0i} - h_{k0,ik}) \quad , \\ R_{ij} &= -\frac{1}{2}(\nabla^2 h_{ij} - h_{00,ij} + h_{kk,ij} - h_{ki,kj} - h_{kj,ki}) \quad , \end{aligned} \quad (21)$$

where time derivatives were neglected, since one assumes that $v \ll c$.

In order to solve the Einstein equations, one first writes the stress-energy tensor for the bumblebee field,

$$T_{B\mu\nu} = \left[\frac{1}{2}b^\alpha b^\beta R_{\alpha\beta} g_{\mu\nu} - b_\mu b^\alpha R_{\alpha\nu} - b_\nu b^\alpha R_{\alpha\mu} \right] \quad , \quad (22)$$

which has a vanishing trace. From the trace of the Einstein equations, one gets

$$R_{\mu\nu} = \kappa \left[T_{\mu\nu} + \frac{1}{2}g_{\mu\nu}T + T_{B\mu\nu} \right] \quad . \quad (23)$$

To get the h_{00} component to first order in the potential U , one writes

$$R_{00} = -\frac{1}{2}(a^2 R_{00} + b^2 R_{11} + 2ab R_{10}) - 2a(a R_{00} + b R_{10}) \quad , \quad (24)$$

which, after a little algebra [11], yields the differential equation

$$\left(\frac{2+5a^2-b^2}{2+5a^2} \right) h_{00,11} + h_{00,22} + h_{00,33} = 0 \quad . \quad (25)$$

This admits the solution

$$h_{00}(x,y,z) = \frac{2M}{\sqrt{c_0^2 x^2 + y^2 + z^2}} \quad , \quad (26)$$

where $c_0^2 = (2+5a^2)/(2+5a^2-b^2)$.

Similarly, the h_{ii} components ($i \neq 1$) obey

$$(2-b^2) h_{ii,11} + 4h_{ii,ii} + 2h_{ii,jj} = (a^2-b^2) h_{00,11} + (2+a^2) h_{00,ii} + a^2 h_{00,jj} \quad . \quad (27)$$

Taking the Ansatz $h_{ii}(x,y,z) = h_{00}(\alpha_1 x, \alpha_2 y, \alpha_3 z)$ and, after some calculation (see Appendix I of Ref. [11]), one can obtain for the coefficients α_i :

$$\alpha_1^2 = 1 - \frac{(2-a^2)b^2}{(2+5a^2)(2-b^2)} \quad , \quad \alpha_i = \alpha_j = 1 \quad . \quad (28)$$

Hence,

$$h_{ii}(x,y,z) = h_{00}(\alpha_1 x, y, z) \equiv \frac{2M}{\sqrt{c_2^2 x^2 + y^2 + z^2}} \quad , \quad (29)$$

with the definition

$$c_2^2 = \alpha_1^2 c_0^2 = \frac{2(2+5a^2-2b^2)-4a^2b^2}{(2+5a^2-b^2)(2-b^2)} \quad . \quad (30)$$

The h_{11} component is now computed; a similar calculation leads to the differential equation [11]

$$h_{11,22} + h_{11,33} = 2M \left[\left(\frac{a^2(c_0^2-1)}{2+3b^2} + c_0^2 \right) \frac{2c_0^2 x^2 - y^2 - z^2}{(c_0^2 x^2 + y^2 + z^2)^{5/2}} - 2c_2^2 \frac{2c_2^2 x^2 - y^2 - z^2}{(c_2^2 x^2 + y^2 + z^2)^{5/2}} \right] \quad , \quad (31)$$

indicating that the solution is a linear combination of h_{00} and h_{22} . Indeed,

$$h_{11}(x,y,z) = - \left(\frac{a^2(c_0^2 - 1)}{2 + 3b^2} + c_0^2 \right) h_{00}(x,y,z) + 2c_2^2 h_{22}(x,y,z) . \quad (32)$$

Proceeding to the off-diagonal component h_{10} , one obtains the differential equation

$$h_{01,22} + h_{01,33} = - \frac{ab}{1 + a^2 + b^2} [h_{00,22} + h_{00,33} + h_{11,22} + h_{11,33} - 2h_{22,11}] . \quad (33)$$

Writing $h_{01} = -ab/(1 + a^2 + b^2)(h_{00} + h_{11} + \delta h_{01})$ leads to

$$\delta h_{01,22} + \delta h_{01,33} = 2h_{22,11} = \frac{4Mc_2^2(2c_2^2x^2 - y^2 - z^2)}{(c_2^2x^2 + y^2 + z^2)^{5/2}} , \quad (34)$$

and hence $\delta h_{01}(x,y,z) = -2c_2^2 h_{22}(x,y,z)$. Therefore,

$$h_{01} = \frac{ab}{1 + a^2 + b^2} \left[\frac{a^2(c_0^2 - 1)}{2 + 3b^2} + c_0^2 \right] h_{00} . \quad (35)$$

Finally, the h_{00} component is computed to second order (see Appendix II of Ref. [11]); it can be shown that only a correction to the first order term $h_{00}^{(1)}$ appears:

$$h_{00} = \frac{2c_0^2[6 + 9b^2 + (15 + 22b^2)a^2] + a^2b^2}{c_0^2(6 + 15a^2 + b^2)(2 + 3b^2)} h_{00}^{(1)} \simeq \left(1 - \frac{b^2}{6} \right) h_{00}^{(1)} . \quad (36)$$

The PPN formalism cannot be straightforwardly used to ascertain its effects, since it relies on a transformation to a quasi-cartesian frame of reference on which, by definition, all diagonal metric components g_{ii} are equal. However, some PPN-like parameters may be extracted from the results, by noticing that

$$\frac{1}{\sqrt{(1 + 2k)x^2 + y^2 + z^2}} \simeq \frac{1}{r} (1 - k \cos^2 \theta) . \quad (37)$$

For h_{00} , one gets

$$h_{00} = \left(1 - \frac{b^2}{6} \right) 2M \frac{1 - (c_0^2 - 1) \cos^2 \theta}{r} . \quad (38)$$

Since no r^{-2} correction appears, the PPN parameter β vanishes in this approach. However, as $h_{11} \neq h_{22} = h_{33}$, the same reasoning allows two parameters analogous to the γ PPN parameter to be obtained: after neglecting the normalization with respect to h_{00} , one gets

$$\begin{aligned} \gamma_1 &= 1 + \cos^2 \theta \times \left[- \left[\frac{a^2(c_0^2 - 1)}{2 + 3b^2} + c_0^2 \right] (1 - c_0^2) + 2c_2^2(1 - c_2^2) \right] \simeq 1 + \frac{b^2}{2} \cos^2 \theta , \\ \gamma_2 &= 1 + (1 - c_2^2) \cos^2 \theta \simeq 1 - \left(\frac{ab}{2} \right)^2 \cos^2 \theta . \end{aligned} \quad (39)$$

As expected, the x-axis LSB produces a stronger effect on the h_{11} component. No clear connection can be derived to link γ with γ_1 and γ_2 , due to the aforementioned anisotropy. However, one can take γ to be of the same order of magnitude as the average of γ_1 and γ_2 , integrated over one orbit:

$$\gamma - 1 \simeq \frac{1}{2}(\gamma_1 + \gamma_2) - 1 \simeq \frac{b^2}{4} \langle \cos^2 \theta \rangle \simeq \frac{b^2}{8} (1 - e^2) , \quad (40)$$

where e is the orbit eccentricity. For low values of e , one gets $\gamma \simeq b^2/8$; The constraint $\gamma = 1 + (2.1 \pm 2.3) \times 10^{-5}$ then enables $|b| \leq 1.9 \times 10^{-2}$.

A discussion concerning the anisotropy of inertia and its effect in the width of resonance lines has been presented as a test between Mach's principle and the Equivalence Principle [21, 22], relying on the hypothetical effect on the proton mass of the proximity to the galactic core. In the present scenario, we note that a radial LSB with the

galactic core acting as source would amount to an axial LSB in a small region such as the Solar System. The bound $\Delta m_P/m_P \leq 3 \times 10^{-22}$, m_P being the proton mass [23], can then be used to obtain

$$\frac{\Delta m_P}{m_P} = \left(1 - \frac{b^2}{6}\right)(c_0^2 - 1) \simeq \frac{b^2}{2} \leq 3 \times 10^{-22} , \quad (41)$$

resulting in the limit $|b| \leq 2.4 \times 10^{-11}$, a much more stringent bound than the obtained above.

5. CONCLUSIONS

In this contribution, the solutions of a gravity model coupled to a vector field where Lorentz symmetry is spontaneously broken are studied, and three different relevant scenarios were highlighted: a purely radial, temporal/radial and temporal/axial LSB.

In the first case, a new black hole solution is found, exhibiting a removable singularity at a horizon of radius $r_s = (2Mr_0^{-L})^{1/(1-L)}$, slightly perturbed with respect to the usual Schwarzschild radius $r_{s0} = 2M$. This has an associated Hawking temperature of $T = (2Mr_0^{-L})^{-L}T_0$, where $T_0 = \hbar/8\pi\kappa_B M$ is the usual Hawking temperature, and protects an intrinsic singularity at $r = 0$. Bounds on deviations from Kepler's law yield $L \leq 2 \times 10^{-9}$.

The temporal/radial scenario produces a slightly perturbed metric that leads to the PPN parameters $\beta \approx 1 - (K + K_r)/M$ and $\gamma \approx 1 - (K + 2K_r)/M$, directly proportional to the strength of the derived effect (given by K and $K_r \sim K$). Since K and K_r are integration constants, no constraints on the physical parameters may be derived from the observed limits on the PPN parameters. Also, an analogy with Rosen's bimetric theory, yields the PPN parameter $\gamma \simeq (A + B)\xi$, ξ being the distance to the central body and A and B parameters ruling the temporal and radial components of the vector field vev .

In the temporal/axial scenario, a breakdown of isotropy is obtained, disallowing a standard PPN analysis. However, the direction-dependent "PPN" parameters $\gamma_1 \simeq b^2 \cos^2 \theta / 2$ and $\gamma_2 \simeq a^2 b^2 \cos^2 \theta / 4$ may be derived, where a and b are respectively the temporal and x -component of the bumblebee vector vev ; naturally, $\gamma_1 \ll \gamma_2$. A crude estimative of the PPN parameter γ yields $\gamma \approx b^2(1 - e^2)/4$, where e is the orbit's eccentricity. Furthermore, a comparison with experiments concerning the anisotropy of inertia produces the bounds $|b| \leq 2.4 \times 10^{-11}$.

REFERENCES

1. "CPT and Lorentz Symmetry II", Ed., V. A. Kostelecký (World Scientific, Singapore, 2002).
2. O. Bertolami, Nucl. Phys. Proc. Suppl. **88**, 49 (2000); O. Bertolami in "Decoherence and Entropy in Complex Systems" (Springer-Verlag, Berlin, 2004).
3. V. A. Kostelecký and S. Samuel, *Phys. Rev.* **D 39**, 683 (1989); *Phys. Rev. Lett.* **66**, 1811 (1991).
4. V. A. Kostelecký and R. Potting, *Phys. Rev.* **D 51**, 3923 (1995).
5. V. A. Kostelecký, *Phys. Rev.* **D 69**, 105009 (2004).
6. R. Bluhm and V. A. Kostelecký, hep-th/0412320.
7. S. M. Carroll, J. A. Harvey, V. A. Kostelecký, C. D. Lane and T. Okamoto, *Phys. Rev. Lett.* **87**, 141601 (2001).
8. O. Bertolami and L. Guisado, *Phys. Rev.* **D 67**, 025001 (2003); *JHEP* **0312**, 013 (2003); O. Bertolami, *Mod. Phys. Lett.* **A 20**, 1359 (2005).
9. V.A. Kostelecký, R. Lehnert and M. J. Perry, *Phys. Rev.* **D 68**, 123511 (2003) ; O. Bertolami, R. Lehnert, R. Potting and A. Ribeiro, *Phys. Rev.* **D 69**, 083513 (2004).
10. H. Sato and T. Tati, *Progr. Theor. Phys.* **47**, 1788 (1972); S. Coleman and S.L. Glashow, *Phys. Lett.* **B 405**, 249 (1997); *Phys. Rev.* **D 59**, 116008 (1999); O. Bertolami and C.S. Carvalho, *Phys. Rev.* **D 61**, 103002 (2000); O. Bertolami, *Gen. Relativity and Gravitation* **34** 707 (2002); R. Lehnert, hep-ph/0312093.
11. O. Bertolami and J. Páramos, *Phys. Rev.* **D 72**, 044001 (2005).
12. R. Grard, M. Novara and G. Scoon, *ESA Bull.* **103**, 11 (2000); L. Iorio, I. Ciufolini and E. C. Pavlis, *Class. Quantum Gravity* **19**, 4301 (2002).
13. S. G. Turyshev *et al.*, gr-qc/0505064.
14. O. Bertolami, J. Páramos and S. G. Turyshev, gr-qc/0601016.
15. M. C. Bento and O. Bertolami, *Phys. Lett.* **B 228**, 348 (1999).
16. E. Fischbach and C.L. Talmadge, "The search for non-Newtonian gravity" (Springer, New York 1999).
17. C. M. Will, *Living Rev. Rel.* **4**, 4 (2001).
18. B. Bertotti, L. Iess and P. Tortora, *Nature* **425**, 374 (2003).
19. N. Rosen, *J. Gen. Rel. and Grav.* **4**, 435 (1973).

20. C.M. Will, "Theory and Experiment in Gravitational Physics", C.M. Will (Cambridge U. P., 1993).
21. S. Weinberg, "Gravitation and Cosmology: Principles and Applications of the General Theory of Relativity" (John Wiley and Sons, New Jersey, 1972).
22. V. A. Kostelecký and C. D. Lane, J. Math. Phys. **40** 6245 (1999).
23. S. K. Lamoreaux, J. P. Jacobs, B. R. Heckel, F. J. Raab, and E. N. Fortson, *Phys. Rev. Lett.* **58**, 746 (1987).